

# High-resolution optical spectroscopy of Tm<sup>3+</sup> ions in LiYF<sub>4</sub>: Crystal-field energies, hyperfine and deformation splittings, and the isotopic structure

Klimin S., Pytalev D., Popova M., Malkin B., Vanyunin M., Korableva S.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

## Abstract

High-resolution optical Fourier spectroscopy was used to study the energies and widths of the crystal-field (CF) levels, hyperfine and deformation level splittings, and isotopic effects in the LiYF<sub>4</sub>: Tm<sup>3+</sup> single crystals. We present the corrected sets of the CF levels and CF parameters for the Tm<sup>3+</sup> ion in LiYF<sub>4</sub>. The observed fine structure of spectral lines is shown to be caused by the hyperfine interaction, random lattice deformations, and isotopic disorder in the lithium sublattice. From a comparison between the observed and calculated fine structures we determine the characteristics of the random lattice deformations in highly diluted activated crystals and obtain an estimate of the fluorine displacements ( $3 \times 10^{-5}$  nm) in the nearest surrounding of the mass defect at the lithium site. © 2010 The American Physical Society.

<http://dx.doi.org/10.1103/PhysRevB.81.045113>

---